

Optimale inhalatie-instructie bij patiënten met longaandoeningen:

de rol van de stichting Inhalatie Medicatie Instructie School (IMIS)

T. Klemmeier-Boekhout

Longverpleegkundige/coördinator stichting IMIS

1. Meek PM, Lareau SS, Fahy BF. Selection of aerosol delivery device. *International Journal of Respiratory Care*, 2006;2:130-133.
2. Rootmensen GN, Van Keimpema ARJ, Jansen HM, De Haan RJ. Predictors of incorrect inhalation technique in patients with asthma or COPD: a study using a validated videotaped scoring method. *Journal of Aerosol Medicine and Pulmonary Drug Delivery*, 2010;23:323-328.
3. Hagedoorn P en Klemmeier-Boekhout T. *Inhalatie technologie & instructie*. Esculaap Media bv, 2014.
4. Hagedoorn P en Klemmeier-Boekhout T. *Zorgatlas Inhalatie-instructie*. Esculaap Media bv, 2014.

COPD, gezondheidsstatus en klinisch relevante verandering

Alma

Referenties

- [1] Westwood M, Bourbeau J, Jones PW, Cerulli A, Capkun-Niggli G, Worthy G. Relationship between FEV1 change and patient-reported outcomes in randomised trials of inhaled bronchodilators for stable COPD: a systematic review. *Respir Res* 2011 Apr 8;12:40-9921-12-40.
- [2] Global Initiative for Chronic Obstructive Lung Disease. *Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease Update 2015*. Available at <http://www.goldcopd.org>. Accessed 01 October 2015.
- [3] NHG-Werkgroep Astma bij volwassenen en COPD. *NHG-Standaard COPD (Derde herziening)*. *Huisarts Wet* 2015;58:108-211
- [4] Jaeschke R, Singer J, Guyatt GH. Measurement of health status. ascertaining the minimal clinically important difference. *Control Clin Trials*. 1989 Dec;10(4):407-15.
- [5] Donohue JF. Minimal clinically important differences in COPD lung function. *COPD* 2005 Mar;2(1):111-124.
- [6] Kocks JW, Tuinenga MG, Uil SM, Berg JW, Ståhl E, Molen T. Health status measurement in COPD: the minimal clinically important difference of the clinical COPD questionnaire. *Respiratory research* 2006;7:62.
- [7] Kon SS, Canavan JL, Jones SE, Nolan CM, Clark AL, Dickson MJ, et al. Minimum clinically important difference for the COPD Assessment Test: a prospective analysis. *Lancet Respir Med* 2014 Mar;2(3):195-203.
- [8] Jones PW. St. George's Respiratory Questionnaire: MCID. *COPD* 2005 Mar;2(1):75-79.

COPD in de eerste lijn; voorspellen van kwaliteit van leven, een proefschriftbespreking

L. Siebeling, Afdeling Huisartsgeneeskunde, Academisch Medisch Centrum, Universiteit van Amsterdam

Referenties

1. Siebeling L, ter Riet G, van der Wal WM, Geskus RB, Zoller M, Muggensturm P, et al. ICE COLD ERIC-- International collaborative effort on chronic obstructive lung disease: exacerbation risk index cohorts-- study protocol for an international COPD cohort study. *BMC Pulm Med* 2009;9:15.
2. Siebeling L, Puhan MA, Muggensturm P, Zoller M, ter Riet G. Characteristics of Dutch and Swiss primary care COPD patients - baseline data of the ICE COLD ERIC study. *Clin Epidemiol* 2011;3:273-83.
3. Siebeling L, Musoro JZ, Geskus RB, Zoller M, Muggensturm P, Frei A, et al. Prediction of COPD-specific health-related quality of life in primary care COPD patients: a prospective cohort study. *NPJ Prim Care Respir Med*. 2014 Aug 28;24:14060.
4. Siebeling L, ter Riet G. Voorspellen van kwaliteit van leven van COPD-patiënten. *Huisarts Wet* 2015;58(10):522-5.
5. Celli BR, Cote CG, Marin JM, Casanova C, Montes de OM, Mendez RA, et al. The body-mass-index, airflow obstruction, dyspnea, and exercise capacity index in chronic obstructive pulmonary disease. *N Engl J Med* 2004 Mar 4;350(10):1005-12.
6. Celli BR, Calverley PM, Rennard SI, Wouters EF, Agusti A, Anthonisen N, et al. Proposal for a multidimensional staging system for chronic obstructive pulmonary disease. *Respir Med* 2005 Dec;99(12):1546-54.
7. Puhan MA, Behnke M, Frey M, Grueter T, Brandli O, Lichtenschopf A, et al. Self-administration and interviewer-administration of the German Chronic Respiratory Questionnaire: instrument development and assessment of validity and reliability in two randomised studies. *Health Qual Life Outcomes* 2004;2:1.
8. Puhan MA, Behnke M, Laschke M, Lichtenschopf A, Brandli O, Guyatt GH, et al. Self-administration and standardisation of the chronic respiratory questionnaire: a randomised trial in three German-speaking countries. *Respir Med* 2004 Apr;98(4):342-50.
9. Schunemann HJ, Puhan M, Goldstein R, Jaeschke R, Guyatt GH. Measurement properties and interpretability of the Chronic respiratory disease questionnaire (CRQ). *COPD* 2005 Mar;2(1):81-9.
10. Bohannon RW. Measurement of Sit-to-Stand Among Older Adults. *Topics in Geriatric Rehabilitation* 2012;28(1):11-16.
11. Jones CJ, Rikli RE, Beam WC. A 30-s chair-stand test as a measure of lower body strength in community-residing older adults. *Res Q Exerc Sport* 1999; Jun;70(2):113-9.
12. Ozalevli S, Ozden A, Itil O, Akkoçlu A. Comparison of the Sit-to-Stand Test with 6 min walk test in patients with chronic obstructive pulmonary disease. *Respir Med* 2007; Feb;101(2):286-93.
13. Puhan MA, Siebeling L, Zoller M, Muggensturm P, ter Riet G. Simple functional performance tests and mortality in COPD. *Eur Respir J* 2013 Oct;42(4):956-63.
14. Lacasse Y, Martin S, Lasserson TJ, Goldstein RS. Meta-analysis of respiratory rehabilitation in chronic obstructive pulmonary disease. A Cochrane systematic review. *Eura Medicophys* 2007 Dec;43(4):475-85.

Toename van de astmaprevalentie in de laatste decennia

Gerrit van Roekel

¹ Bindels PJE, Van de Griendt EJ et al; NHG-Standaard Astma bij kinderen; *Huisarts Wet* 2014;57(2):70-80

² Pouwelse M. et al; Astma bij kinderen: hoe zeker is de diagnose? *huisarts & wetenschap*, 2015 58(8) augustus

³ van Schayck CP, Smit HA; The prevalence of asthma in children: a reversing trend; *Eur Respir J* 2005; 26: 647–650

⁴ Anandan C, Nurmatov U, van Schayck O C P, Sheikh A; Is the prevalence of asthma declining? Systematic review of epidemiological studies. *Allergy* 65 (2010) 152–167 © 2009 John Wiley & Sons A/S 152

⁵ Asher MI, Montefort S, Björkstén B, Lai CK, Strachan DP, Weiland SK; Worldwide time trends in the prevalence of symptoms of asthma, allergic rhinoconjunctivitis, and eczema in childhood; *Lancet* 2006; 368: 733–43

⁶ To T, Stanojevic S, Moores G, et al. Global asthma prevalence in adults: findings from the cross-sectional world health survey. *BMC Public Health* 2012; 12:204.

⁷ de Nijs SB, Venekamp LN and Bel EH; Adult-onset asthma: is it really different? *Eur Respir Rev* 2013; 22: 127,