

**Referenties bij het artikel 'Inspanningsgeïnduceerde luchtwegvernauwing en het onderscheid met inspanningsastma
door Gerrit van Roekel, kaderhuisarts astma/COPD**

[¹] Storms W., Review of exercise-induced asthma, *Medicine & Science in sports & exercise*, 2003 the american college of sports medicine, 1464-1469

[²] Parsons JP, Mastrorade JG, exercise-induced bronchoconstriction in athletes, *Chest* 2005; 128; 3966-3974

[³] Billen A, Dupont L. Exercise induced bronchoconstriction and sports, *postgrade Med.J.* 2008; 84; 512-517

[⁴] Billen A, Dupont L. Exercise induced bronchoconstriction and sports, *postgrade Med.J.* 2008; 84; 512-517

[⁵] Anderson SD, How does exercise cause asthma attacks? *Current Opinion in Allergy and Clinical Immunology* 2006, 6; 37-42

[⁶] Weiss P, Rundell KW, Imitators of exercise-induced bronchoconstriction, *Allergy, Asthma & Clinical Immunology*, 2009, 5:7

[⁷] Anderson SD, Argyros GJ, Magnussen H, Holzer K, Provocation by eucapnic voluntary hyperpnoea to identify exercise induced bronchoconstriction, *Br J Sports Med* 2001; 35: 344-347

[⁸] Hull JH, Hull PJ, Parsons JP, Dickinson, JW, Ansley L, Approach to the diagnosis and management of suspected exercise-induced bronchoconstriction by primary care physicians, *BMC Pulmonary medicine* 2009, 9:29

[⁹] Kelly KD, Spooner CH, Rowe BH, *Eur Respir J* 2001; 17: 39-45